



PrimeraOne®

Imidacloprid 75WSP INSECTICIDE

For foliar and systemic insect control on turfgrass, nursery, greenhouse, commercial and residential landscape ornamentals, on fruit and nut trees, and on Ornamental and vegetable plants in greenhouses, nurseries and interior landscapes.

ACTIVE INGREDIENT:

Imidacloprid, 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....75%

OTHER INGREDIENTS:.....25%

Total:100%

This product contains 0.075 lbs (1.2 oz) (34.02 grams) imidacloprid per packet.

Keep water-soluble packets in this container and store in a cool dry place but not below freezing (32°F).

DO NOT remove packets from container except for immediate use.

EPA Reg. No.: 83100-26-88975 EPA Est. No.: 33967-NJ-01

KEEP OUT OF REACH OF CHILDREN CAUTION

PRODUCT OF CHINA

PRECAUCION AL USUARIO: Si usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

Manufactured for:
PrimeraTurf Inc.
3677 Guildhall Trail
Marietta, GA 30066

FOR ADDITIONAL PRECAUTIONARY, HANDLING, AND USE STATEMENTS, SEE INSIDE OF THIS BOOKLET

**Water Soluble Packaging
Net Contents: 1.6 ounces**

TABLE OF CONTENTS

	page
PRECAUTIONARY STATEMENTS	1
FIRST AID	1
ENVIRONMENTAL HAZARDS	1
DIRECTIONS FOR USE	1
AGRICULTURAL USE REQUIREMENTS	2
NON-AGRICULTURAL USE REQUIREMENTS	2
PRECAUTIONS FOR AQUATIC AREAS	2
Runoff Management.....	2
Endangered Species Notice.....	2
Resistance Management.....	2
Spray Drift Management.....	2
Use Information.....	3
Use Restrictions.....	3
ROTATION CROP RESTRICTIONS	3
PREPARATION OF TANK MIXES	4
CHEMIGATION	4
Safety Devices for Systems Connected to a Public Water Supply.....	4
Safety Devices for Systems Not Connected to a Public Water Supply.....	5
APPLICATIONS	5
Turf Grass.....	5
Landscape Ornamentals.....	5
Pome Fruits.....	7
Pecans.....	7
Grapes.....	8
Citrus.....	8
Avocado.....	8
Grassy Areas In Nurseries, Nursery, And Greenhouse Grown Ornamentals.....	9
STORAGE AND DISPOSAL	11
CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY	12

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled, or absorbed through skin. Causes eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust or vapor. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Keep children or pets off treated area until spray is dry.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and Other Handlers Must Wear:

- Long-sleeved shirt and long pants
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton.
- Shoes plus socks

Follow manufacturers' instructions for cleaning/maintaining personal protective equipment, PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering controls statements:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

IF SWALLOWED

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by a poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

IF ON SKIN

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15 to 20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

In case of emergency call your local poison control center for assistance. Have a product container or label with you when calling a poison control center or doctor, or going for treatment.

Note To Physician: No specific antidote is available. Treat the patient symptomatically.

ENVIRONMENTAL HAZARDS

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Do not formulate this product into other end-use products.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinyl chloride (PVC) or viton.
- Shoes plus socks

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep children and pets off treated area until dry.

OBSERVE THE FOLLOWING PRECAUTIONS WHEN MIXING AND APPLYING IN THE VICINITY OF AQUATIC AREAS SUCH AS LAKES, RESERVOIRS, RIVERS, PERMANENT STREAMS, MARSHES OR NATURAL PONDS, ESTUARIES AND COMMERCIAL FISH PONDS.

RUNOFF MANAGEMENT

DO NOT cultivate within 10 feet of the aquatic areas to allow growth of vegetative filter strip. When used on erodible soils, use best management practices for minimizing runoff. Consult your local Natural Resources Conservation Service for recommendations in your use area.

ENDANGERED SPECIES NOTICE

Under the Endangered Species Act, it is a Federal Offense to use any pesticide in a manner that results in the death of a member of an endangered species. Consult your local county bulletin, County Extension Agent, or Pesticide State Lead Agency for information concerning endangered species in your area.

RESISTANCE MANAGEMENT

Certain insects may develop resistance to insecticides after repeated use. Use different resistance management practices such as rotating classes of insecticides to help delay or minimize insect resistance.

This product contains the active ingredient imidacloprid, which is a Group 4A insecticide. Repeated use of Group 4A insecticides may lead to insect pests that become resistant to imidacloprid or other neonicotinoids (Group 4A) insecticides.

To reduce the chances of development of resistance to Group 4A insecticides, do not make more than three (3) consecutive foliar applications of this product and/or other Group 4A insecticides with similar modes of action. In addition, PrimeraTurf Inc. strongly recommends the use of other insecticides with a different mode of action prior to or after application of this product. This strategy of insecticide rotation in concert with other IPM practices is considered an effective way to delay or minimize an insect's ability to develop resistance to this class of chemistry.

Some Group 4A neonicotinoid products used as foliar treatments include the active ingredients thiamethoxam (found in Actara® and Centric®), acetamiprid (found in Assail® and Intruder™), thiacloprid (found in Calypso®), and imidacloprid (found in Leverage®, Provado®, and Trimax™). Some 4A Group neonicotinoid products used as soil treatment include thiamethoxam (found in Platinum®) and imidacloprid (found in Admire®).

Additional information on insect resistance management may be obtained from your local extension specialist, certified crop advisor and/or product manufacturer, or from the Insecticide Resistance Action Committee (IRAC) on the web at <http://irac-online.org/>.

SPRAY DRIFT MANAGEMENT

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all of these factors when making application decisions. Avoiding spray drift is the responsibility of the applicator.

Mixing and Loading Requirements

To avoid potential contamination of groundwater, use a properly designed and maintained containment pad for mixing and loading of any pesticide into application equipment. If containment pad is not used, maintain a minimum distance of 25 feet between mixing and loading areas and potential surface to groundwater conduits such as field sumps, uncased well heads, sinkholes, or field drains.

Importance of Droplet Size

An important factor influencing drift is droplet size. Small droplets (<150-200 microns) drift to a greater extent than large droplets. Within typical equipment specifications, make applications to deliver the largest droplet spectrum that provides sufficient control and coverage. Formation of very small droplets may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray boom pressure.

Release the spray at the lowest possible height consistent with good pest control and flight safety. DO NOT make applications more than 10 feet above the crop canopy.

Wind Speed Restrictions

Drift potential increases at wind speeds of less than 3 mph (due to inversion potential) or more than 10 mph. However, many factors, including droplet size, canopy and equipment specifications determine drift potential at any given wind speed. DO NOT apply when winds are greater than 15 mph and avoid gusty and windless conditions. Risk of exposure to sensitive aquatic areas can be reduced by avoiding applications when wind direction is toward the aquatic area.

Restrictions During Temperature Inversions

Because the potential for spray drift is high during temperature inversions, DO NOT make ground applications during temperature inversions. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain close to the ground and move laterally in a concentrated cloud. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the morning. Their presence can be indicated by ground fog; however if fog is not present, inversions can also be identified by the movement of smoke from a ground source. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical mixing.

Airblast (Air Assist)

Airblast sprayers carry droplets into the canopy of trees/vines via a radially, or laterally directed air stream. Follow the specified drift management practices:

- Adjust deflectors and aiming devices so that spray is only directed into the canopy;
- Block off upward pointed nozzles when there is no overhanging canopy;
- Use only enough air volume to penetrate the canopy and provide good coverage;
- DO NOT allow the spray to go beyond the edge of the cultivated area (i.e. turn off sprayer when turning at end rows);
- Only spray inward, toward the orchard or vineyard, for applications to the outside rows.

No-Spray Zone Requirements for Foliar Applications

DO NOT apply by ground within 25 feet lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, estuaries and commercial fish farm ponds.

USE INFORMATION

Thorough uniform coverage is necessary to achieve optimal control. A spray adjuvant may be used to improve coverage.

This product may not knockdown established and heavy insect populations. Two applications may be required to achieve control; retreat if needed and as directed on this label.

Tank mix this product with other insecticides as recommended for knockdown of pests or for improved control of other pests.

USE RESTRICTIONS (All Uses)

- DO NOT make a foliar application of any chloronicotinyl insecticide for resistance management purposes following a soil application of this product on the same crop.
- DO NOT use product packets in a tank mix with products that contain boron or release free chlorine; the PVA packet reacts with boron or free chlorine to produce a plastic that is not soluble in water. NOTE: Normal chlorinated water is acceptable for mixing.
- DO NOT use this product on commercial sod farms.
- DO NOT allow livestock to graze in treated areas or use clippings from treated areas for feed or forage unless specified otherwise on this label.
- DO NOT apply this product to soils that are waterlogged or saturated.
- DO NOT allow runoff or puddling of irrigation water following application.
- DO NOT allow leachate to run off for the first 10 days after application or reduced efficacy may result.
- DO NOT exceed the total 5.375 packets (8.6 oz) of this product (0.4 lbs AI) / Acre per year specified for the uses indicated on this label.

ROTATION CROPS RESTRICTIONS

Crops which are listed on imidacloprid labels or crops that have existing tolerances for imidacloprid may be planted in treated areas as soon as practical after the last imidacloprid application. Crops that are not found on an imidacloprid label, or crops that do not have existing tolerances for imidacloprid, may not be planted in treated areas for 12-months after the last application.

Refer to the table below for plantback intervals for different crops. Note that if cover crops are planted any time after an application of this product, those crops may not be grazed or harvested for food or feed.

Crops	Plantback Interval
All crops on this label, plus the following crops not on this label: barley, canola, corn (field, sweet and pop), rapeseed, sorghum, sugar beet, and wheat	No restrictions
Cereals (including buckwheat, millet, oats, rice, rye, and triticale), soybeans and safflower	30 days
Onion and bulb vegetables	10 months
All other crops	12 months

PREPARATION OF TANK MIXES

Handling PVA pouches: The enclosed packets containing this product are water-soluble and will completely dissolve in water. To prevent the packets from dissolving before they are added to the spray tank, do not get the packets wet from handling with wet hands or wet gloves. The packets can break open if handled roughly. Remove the packets from the outer bag but do not open the packets. Keep unused packets sealed in the outer bag. To prepare a tank mix of this product, follow the steps below. If tank mixing this product with other components, read the section on Compatibility, below.

1. Fill the spray tank with 1/4 to 1/2 of the required amount of water and begin agitation.
2. Add the specified amount of this product in PVA packets and allow packets to fully dissolve.
3. Fill the tank with the remaining water needed. Maintain sufficient agitation during mixing and application.

If this product is to be tank mixed with other pesticides and/or fertilizer solutions, check the compatibility (refer to the Compatibility section below) before preparing tank mixes.

To prepare tank mixes of this product with other pesticides, use the following order of mixing:

1. PrimeraOne Imidacloprid 75WSP PVA packets;
2. Other wettable powders or wettable granules;
3. Flowables or suspension concentrates;
4. Emulsifiable concentrates.

Agitate the solution as each component is added. Add the next component only after the previous one is thoroughly mixed. If needed, add a compatibility agent when adding a fertilizer solution to the mix. Add the remaining quantity of water as the final step. To ensure a uniform spray mixture, maintain constant agitation during both mixing and application.

Compatibility PVA packets that are tank mixed with products that contain boron or that release free chlorine will react to form a plastic that is insoluble in water or solvents such as alcohol, kerosene, diesel oils or gasoline. Further information is available from your local PrimeraTurf Inc. representative. Conduct the following test for compatibility of the intended tank mix partner product(s) before adding this product to the spray or mix tank:

1. In a pint or quart jar, add proportionate amounts of each tank mix component in the order provided in the directions above.
2. Replace the cap, shake for 5 minutes, and allow the mixture to settle for 5 minutes.
3. Observe the jar for signs indicating an incompatible mixture. If the contents can be re-mixed by shaking and readily re-suspends, it is considered compatible. If the mixture separates out, foams, or forms a gel or lumps, then the mixture is not compatible.

APPLICATION THROUGH IRRIGATION SYSTEMS (CHEMIGATION)

Apply this product at rates specified on this label either alone or in tank mixture with other pesticides and chemicals registered for application through irrigation systems. The normal dilution ratio is 1:10 to 1:200, depending on the system. Always meter the product into the irrigation water during the first part of the irrigation cycle. Mix the product separately prior to injection. Agitate as necessary if the mixture is allowed to stand more than 24 hours.

- DO NOT connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.
- Apply this product only through micro-irrigation (individual spaghetti tube), drip irrigation, overhead irrigation, and ebb and flood or hand-held or motorized calibrated irrigation equipment. DO NOT apply this product through any other type of irrigation system. Crop injury or lack of effectiveness can result from non-uniform distribution of treated water.
- Be sure to remove scale, pesticide residue and other foreign matter from the tank and entire irrigation system prior to application.
- A person knowledgeable of the chemigation system and responsible for its operation, or a person who is under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
- If you have any questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

SAFETY DEVICES FOR IRRIGATION SYSTEMS CONNECTED TO PUBLIC WATER SUPPLIES:

If the source of water for your irrigation system is a public water supply, follow the instructions below.

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.
3. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
4. The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
5. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. DO NOT apply when wind speed favors drift beyond the area intended for treatment.

SAFETY DEVICES FOR IRRIGATION SYSTEMS NOT CONNECTED TO A PUBLIC WATER SUPPLY:

1. The system must contain a functional check valve, vacuum relief valve and low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where the pesticide distribution is adversely affected.
6. Systems must use a metering pump such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. DO NOT apply when wind speed favors drift beyond the area intended for treatment.

TURFGRASS APPLICATIONS

PrimeraOne Imidacloprid 75 WSP Insecticide can be used for the control of listed soil inhabiting pests as directed on turfgrass.

The active ingredient in PrimeraOne Imidacloprid 75 WSP Insecticide has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods.

TURFGRASS APPLICATIONS: around airports, athletic fields, cemeteries, golf courses, homes and multi-family residential buildings, office buildings and office parks, public parks and playground areas, and shopping centers	
PEST	RATE: PrimeraOne Imidacloprid 75WSP
Annual bluegrass weevil Asiatic garden beetle Billbug Black turfgrass ataeenius European chafer European Crane Fly Green June beetle Japanese beetle May or June Beetles Northern masked chafer Oriental beetle Phyllophaga spp. Southern masked chafer	1 packet (1 .6 oz) / 8,250 - 11 ,000 ft ² or 4 - 5.375 packets (6.4 - 8.6 oz) / Acre
For Suppression of: Chinchbug ² Cutworms For Control of: Mole Crickets ¹	1 packet (1 .6 oz) / 8,250 ft ² or 5.375 packets (8.6 oz) / Acre
<p>Application Instructions: The use of accurately calibrated equipment normally used for the application of turfgrass insecticides is required. Use equipment which will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off target drift. Check calibration periodically to ensure that equipment is working properly.</p> <p>Make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch. Apply this product in sufficient water to provide adequate distribution in the treated area. Rainfall or irrigation must occur within 24 hours of application to move this product vertically through the thatch and into the soil.</p> <p>Wait until after sufficient rainfall or irrigation has occurred to mow the grass.</p> <p>Remarks Annual Bluegrass weevil, Billbugs, European Crane Fly, and Grubs: For best results, make applications before egg hatch. ¹Mole Crickets: Make applications before or during the peak egg hatching period, this product may be applied with acurative insecticide when adults or large nymphs are present and actively tunneling, ²Chinchbugs: Make applications before hatching of first instar nymphs.</p> <p>RESTRICTIONS:</p> <ul style="list-style-type: none"> • Do not apply more than 5.375 packets (8.6 oz) (0.4 lb of active ingredient) per acre per year. • DO NOT make applications when turfgrass areas are waterlogged or the soil is saturated with water. Adequate distribution of the active ingredient cannot be achieved when these conditions exist. The treated turf area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile. 	

APPLICATION TO LANDSCAPE ORNAMENTALS

Use this product as directed below on ornamentals in and around commercial and residential landscapes and interior plantscapes to control or suppress listed insects.

This is a systemic product and will be translocated upward into the plant system from root uptake. Plants absorb this product from either foliar or soil applications. Apply this product to areas where the growing portion of the target plant can absorb the active ingredient. The addition of a nitrogen containing fertilizer, where applicable, into the solution has been shown to enhance the uptake of the active ingredient. Application can be made by foliar application or soil applications; including soil injection, drenches, and broadcast sprays. Foliar applications offer locally systemic activity against insect pests.

Woody Perennials: When making soil applications to plants with woody stems, systemic activity will be delayed until the active ingredient is translocated throughout the plant. In some cases, this translocation delay could take 60 days or longer. For this reason, applications must be made prior to anticipated pest infestation to achieve control.

Bark Media: Treatments of this product to media with 30 - 50% or more bark content may confer a shorter period of protection.

Ant Management Programs: Use this product to control aphids, scale insects, mealybugs and other sucking pests on ornamentals to limit the honeydew available as a food source for ant populations. Applications can then be supplemented with residual sprays, bait placements or other ant control tactics to further reduce the pest population.

ORNAMENTAL TREES, SHRUBS, EVERGREENS, FLOWERS, FOLIAGE PLANTS, GROUNDCOVERS, INTERIOR PLANTSCAPES, NON-BEARING FRUIT & NUT TREES, VEGETABLE PLANTS (intended for Resale only) (in and around the perimeter of industrial and commercial buildings and residential areas, and state, national, and private wooded and forested areas for the insect pests listed below)

Foliar Applications

PEST	RATE: PrimeraOne Imidacloprid 75WSP
Adelgids Aphids Japanese beetles (adult) Lace bugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassywinged sharpshooter) Leafminers Mealybugs Sawfly larvae Thrips (suppression) Whiteflies	1 packet (1.6 oz) in 300 gal. of water

Foliar Application Instructions
Apply this product in a sufficient volume of water to uniformly cover the treatment area. Foliar applications will provide systemic activity against target pests.
If plants (such as holly, pine or ivy) have foliage that is difficult to wet, PrimeraTurf Inc. recommends this product be applied with a spreader/sticker.
Time applications to occur before heavy pest populations arise; make repeat applications as necessary.

Soil (Broadcast) Applications

Target Pests	Rate
White grub larvae (including Asiatic garden beetle, chafer, <i>Phyllophaga</i> spp., Japanese beetle larvae, and Oriental beetle)	1 packet (1.6 oz) / 8,250 - 11,000 ft ² or 4 - 5.375 packets (6.4 - 8.6 oz) / Acre

Soil Application Instructions
Mix the specified amount of this product in a sufficient volume of water to uniformly cover the treatment area. Apply in a minimum of 2 gallons of water per 1,000 sq. ft. After application, irrigate the treated areas to incorporate this product into the soil.

Restrictions:

DO NOT apply more than 5.375 packets (8.6 oz). (0.4 lb active ingredient) per acre per year by broadcast application.

ORNAMENTAL TREES, SHRUBS, FLOWERS AND GROUNDCOVERS (in and around the perimeter of industrial and commercial buildings and residential areas, and state, national, and private wooded and forested areas for the insect pests listed below)

For control of: Adelgids, Alder borer, Aphids, Asian longhorned beetle, Black vine weevil larvae, Bronze birch borer, Emerald ash borer, Eucalyptus longhorned borer, Flat-headed borers (including Bronze birch and Alder), Japanese beetles, Lace bugs, Leaf beetles (including elm and viburnum leaf beetles), Leafhoppers (including glassy-winged sharpshooter), Leafminers, Mealybugs, Pine tip moth larvae, Psyllids, Royal palm bugs, Sawfly larvae*, Soft scales, White grub larvae, Whiteflies
For suppression of: Armored scales, Thrips

USE RATE	APPLICATION SITE
	TREES
Diameter at Breast Height (DBH) is measured at 4.5 feet from the ground. Use the following rates as a function of tree diameter at breast height (DBH): Apply 1 packet (1.6 oz) per 12 – 48 inches of trunk diameter (DBH). You may only use the higher rate on trees >15 inches (DBH) to control: Asian longhorned beetle, Emerald ash borer, Eucalyptus longhorned borer, Bronze birch borer, and Alder borer. To calculate the higher rates, divide trunk diameter by 12 - 23 inches. Refer to example calculations below. RESTRICTION: DO NOT apply more than 5.375 packets (8.6 oz) (0.4 lb AI) per acre per year.	Soil Injection: GRID SYSTEM: Holes must be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per tree. NEW YORK SPECIFIC RESTRICTION: No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York. Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone. For Control of Specified Borers: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.

EXAMPLE CALCULATIONS:

Example 1 (to calculate the standard rate): If you have three trees having DBH of 8, 10 & 16 inches, the total cumulative inches of trunk diameter is 34 inches (8 + 10 + 16 = 34)
34/48 = .708x1.6fl. oz (1 packet) = 1.13oz -OR- 34/24 = 1.417x1.6fl. oz. (1 packet) = 2.27 oz
If you have a single tree with a DBH of 12 inches, the lower rate range will be:
12/48 = 0.25 x 1.6 oz (1 packet) = 0.4 oz -OR- 12/24 = 0.5 x 1.6 oz (1 packet) = 0.8 oz
Example 2 (To calculate the higher rate): If you have three trees having DBH of 15, 20 & 25 inches, the total cumulative inches of trunk diameter is 60 inches (15 + 20 + 25 = 60)
60/23 = .26 x 1.6 fl. oz (1 packet) = 4.17 oz -OR- 60/12 = 5 x 1.6 fl. oz. (1 packet) = 8 oz
If you have a single tree with a DBH of 30 inches, the higher rate range will be:
30/23 = 1.3x1.6oz(1 packet) = 2.08 oz -OR- 30/12 = 2.5 x 1.6 oz (1 packet) = 4.0 oz

SHRUBS

1 packet (1.6 oz) per 24 - 48 feet cumulative shrub height	<p>Soil Injection: Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per shrub.</p> <p>NEW YORK SPECIFIC RESTRICTION: No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York.</p> <p>Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.</p>
--	--

FLOWERS & GROUNDCOVERS

1 packet (1.6 oz) / 8,250 - 11,000 ft ² or 4 - 5.375 packets (6.4 - 8.6 oz) /Acre	Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. Irrigate immediately following application to established plants.
--	---

Remarks
*Sawfly larvae feed on mature foliage beginning in early spring. Make treatments in the fall before pine sawfly emergence in spring to allow adequate time for imidacloprid translocation into mature foliage.
Diameter at Breast Height (D.B.H.) is measured at 4.5 feet from the ground.

Restrictions
DO NOT apply more than 5.375 packets (8.6 ounces) (0.4 lbs AI) per acre per year.
DO NOT harvest or consume fruits or nuts from trees that have been treated within 1 year of application.

APPLICATION TO POME FRUITS

POME FRUITS: Apple, Crabapple, Loquat, Mayhaw, Pear, Pear (oriental), Quince (around perimeter of industrial and commercial buildings and on residential areas)

PEST	USE RATE
Aphids (except Woolly apple aphid) Leafhoppers (including glassy-winged sharpshooter) Leafminer Mealybugs* San Jose scale*	1 packet (1.6 oz) in 300 Gallons of Water 1.3125 packets (2.1 oz) / Acre ¹

Remarks
Apply specified dosage as foliar spray as needed after petal-fall is complete.
For control of Rosy apple aphid, apply prior to leafrolling caused by the pest.
For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. This product will not control late stage larvae.
For San Jose Scale, time applications to the crawler stage. Treat each generation.
For late season (preharvest) control of leafhopper species, apply this product while most leafhoppers are in the nymphal stage.
For control of mealybugs, ensure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybugs.
¹The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.

Restrictions
DO NOT apply more than 1.3125 packets (2.1 oz) per acre in a single application.
DO NOT make more than 5 applications per year.
Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.
* Not for use in California for control on pears.

APPLICATION TO PECANS

PECANS* (around perimeter of industrial and commercial buildings and on residential areas)

PEST	USE RATE
Yellow pecan aphid Black margined aphid Pecan leaf phylloxera Pecan spittlebug Pecan stem phylloxera	1 packet (1.6 oz) in 300 gallons of water 1.3125 packets (2.1 oz) / Acre ¹

Remarks

Make foliar applications as pests begin to build before populations become extreme. Two applications at a 10 to 14-day interval may be required to achieve control. Scout and retreat if needed. Thorough uniform coverage of foliage is necessary for control. Addition of an organosilicone-based spray adjuvant at a rate not to exceed the adjuvant manufacturer's specified use rate may improve coverage.

¹The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.

Restrictions

DO NOT apply more than a total of 3.9375 packets (6.3 oz) of this product per acre per year.

DO NOT make more than 3 applications per year.

Allow 10 or more days between applications.

* Use on pecans not permitted in California unless otherwise directed by specific supplemental labeling.

APPLICATION TO GRAPES

GRAPES: (around perimeter of industrial and commercial buildings and on residential areas)

PEST	RATE
Leafhoppers (including glassy-winged sharpshooter) Mealybugs	1 packet (1.6 oz) in 300 gallons of water 1.0 oz / Acre ¹
Remarks ¹ Apply specified dosage as a foliar spray using 200 gallons of water per acre.	
Restrictions DO NOT apply more than a total of 1.25 packets (2.0 oz) of this product per acre per year. Allow at least 14 days between applications. Applications may be applied up to and including day of harvest.	

APPLICATION TO CITRUS

CITRUS: Citrus and Citrus hybrids, Orange (sweet and sour), Calamondin, Grapefruit, Kumquat, Lemon, Lime, Pummelo, Tangerine, Tangelo (around perimeter of industrial and commercial buildings and on residential areas)

PEST	USE RATE
Aphids Asian citrus psyllid Black fly Citrus leafminer Leafhoppers/Sharpshooters Mealybugs Scales Termites (FL only) Whiteflies	1 packet (1.6 oz) in 300 gallons of water 1.3125 packets (2.1 oz)/Acre ¹
Remarks Apply specified dosage as foliar spray as needed after petal-fall is complete. For first generation leafminer control, make first application as soon as petal-fall is complete. Greatest leafminer control will result from the earliest possible application. For second and succeeding generations of leafminer, optimal control is obtained from applications made early in the adult flight against egg and early instar larvae. A second application may be required 10 days later if severe pressure continues or if generations are overlapping. A single application may result in suppression only. This product will not control late stage larvae. For late season (preharvest) control of leafhopper species, apply this product while most leafhoppers are in the nymphal stage. For control of mealybugs, ensure good spray coverage of the trunk and scaffolding limbs or other resting sites of the mealybugs. ¹ The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.	
Restrictions DO NOT apply more than 1.3125 packets (2.1 oz) per acre in a single application. DO NOT make more than 5 applications per year. Allow 10 or more days between applications. Allow at least 7 days between last application and harvest.	

APPLICATION TO AVOCADO

AVOCADO: (around perimeter of industrial and commercial buildings and on residential areas)

PEST	RATE
Aphids Avocado lacebug Leafhoppers Whiteflies	1 packet (1.6 oz) in 300 gallons of water 1.3125 packets (2.1 oz) / Acre ¹
Remarks ¹ The amount of this product required per acre will depend on tree size and volume of foliage present. The rate per acre is based on a standard of 400 gallons of dilute spray solution per acre for large trees.	
Restrictions DO NOT apply more than a total of 1.3125 packets (2.1 oz) of this product per acre per year. Allow at least 14 days between applications. Allow at least 7 days between application and harvest.	

APPLICATION TO GRASSY AREAS IN NURSERIES, NURSERY, AND GREENHOUSE GROWN ORNAMENTALS

Use this product on grassy areas in nurseries, around and on nursery grown ornamentals, and in planting rows in nurseries to control listed pests. Make application prior to anticipated pest infestation to maximize control. Rainfall, irrigation and mechanical incorporation after application will aid in maximizing control.

The active ingredient in this product has sufficient residual activity so that applications can be made preceding the egg laying activity of the target pests. High levels of control can be achieved when applications are made preceding or during the egg laying period. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trapping or other methods.

GRASSY AREAS (in Field & Forest Nurseries)	
PEST	DOSAGE
For control of: Annual bluegrass weevil Asiatic garden beetles (<i>Maladera</i> spp.) Billbugs Black turfgrass ataenius European chafer European Crane Fly Green June beetle Japanese beetle May or June beetles Northern masked chafer Oriental beetle (<i>Phyllophaga</i> spp.) Southern masked chafer	1 packet (1 .6 oz) / 8,250 - 11 ,000 ft ² or 4 - 5.375 packets (6.4 - 8.6 oz) / Acre
For control of: Mole crickets For suppression of: Chinchbugs Cutworms	1 packet (1. 6 oz)/ 8,250 ft ² or 5.375 packets (8.6 oz) / Acre
<p>Application Instructions Apply this product in sufficient water to provide adequate distribution in the treated area. The use of accurately calibrated equipment normally used for soil application of insecticides is required. Use equipment that will produce a uniform, coarse droplet spray, using a low pressure setting to eliminate off-target drift. Check calibration periodically to ensure that equipment is working properly.</p> <p>Remarks For control of grubs, billbugs, European crane fly and annual bluegrass weevil, make application prior to egg hatch of the target pest. For control of mole crickets make application prior to or during the peak egg hatch period. When adults or large nymphs are present and actively tunneling, this product should be accompanied by a curative insecticide. For suppression of chinchbugs, make application prior to the hatching of the first instar nymphs. Consult your local State Agricultural Experiment Station, or State Extension Turf Specialists for more specific information regarding timing of application. Mowing of the vegetation in the area to be treated to a height of 3 inches or less prior to application will improve the consistency of control. NOTE: Irrigation or rainfall is needed within 24 hours after application to move the active ingredient through the thatch.</p> <p>Restrictions DO NOT apply more than 5.375 packets (8.6 oz). (0.4 lb of active ingredient) per acre per year. DO NOT mow grassy area until after irrigation or rainfall has occurred so that uniformity of application will not be affected. DO NOT allow runoff or puddling of irrigation water following application. DO NOT apply this product to water-logged or saturated areas. Application of this product to water-logged or saturated areas will not allow penetration into the root zone of the plant. DO NOT graze treated areas or use clippings from treated areas for feed or forage.</p>	

ORNAMENTAL TREES (including non-bearing fruit & nut trees), SHRUBS, EVERGREENS, FLOWERS, FOLIAGE PLANTS, GROUNDCOVERS, INTERIOR PLANTSCAPES, VEGETABLE PLANTS* (on and around field-grown nursery and container stock, indoor and outdoor ornamentals (including both greenhouse and interior plantscapes) and on ornamentals grown in flats, benches or beds)

Foliar Applications	
PEST	DOSAGE
For control of: Adelgids Aphids Japanese beetle (adult) Lacebugs Leaf beetles (including elm and viburnum leaf beetles) Leafhoppers (including glassy-winged sharpshooter) Leafminers Mealybugs Sawfly larvae Whiteflies For suppression of: Thrips	1 packet (1.6 oz) in 300 gal. of water
<p>Foliar Application Instructions Apply this product in a sufficient volume of water to uniformly cover the treatment area. Foliar applications will provide systemic activity against target pests. If plants (such as holly, pine or ivy) have foliage that is difficult to wet, PrimeraTurf Inc. recommends this product be applied with a spreader/sticker. Time applications to occur before heavy pest populations arise; make repeat applications as necessary.</p>	

Soil (Broadcast) Applications

PEST	DOSAGE
For control of: White grub larvae (including Japanese beetle, Masked Chafers, European Chafer, Oriental beetle, Asiatic Garden beetle)	1 packet (1.6 oz) / 8,250 - 11 ,000 ft ² or 4 - 5.375 packets (6.4 - 8.6 oz) / Acre
<p>Soil (Broadcast) Application Methods Mix the required amount of this product in sufficient water to uniformly and accurately cover the area being treated. DO NOT use less than 2 gallons of water per 1 000 sq. ft. of treatment area. Irrigate thoroughly to incorporate the product into the upper soil profile. Incorporate application into the soil before planting or apply after plants are established. For applications made to established plants, irrigate thoroughly after application.</p> <p>Remarks Mowing of the vegetation in the area to be treated to a height of 3 inches or less prior to application will improve the consistency of control.</p> <p>Restrictions DO NOT apply more than 5.375 packets (8.6 oz) (0.4 lb of active ingredient) per acre per year. DO NOT allow runoff or puddling of irrigation water following application. DO NOT apply this product to water-logged or saturated areas. Application of this product to water-logged or saturated areas will not allow penetration into the root zone of the plant. DO NOT exceed an application frequency of more than once each 16 weeks on nursery ornamentals with a production cycle of less than one (1) year. DO NOT exceed one (1) application per year on nursery ornamentals with a production cycle of greater than one year. *Only for use on vegetable plants intended for resale including: Broccoli, Chinese broccoli, Broccoli raab, Brussels sprouts, Cabbage, Chinese Cabbage, Cauliflower, Collards, Eggplant, Ground cherry, Kale, Kohlrabi, Lettuce, Mustard greens, Pepinos, Peppers, Potatoes, Rape greens, Sorghum, Sugarbeets, Tomatillo, and Tomato. DO NOT make a foliar application of this product following a soil application in the same crop for resistance management purposes.</p> <p>Rotational Crops: Food Crops: Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on any imidacloprid label, or for crops for which no tolerances for the active ingredient have been established, a 12 month plant-back interval must be observed.</p>	

SOIL INJECTION & BASAL DRENCH APPLICATIONS: NURSERY, GREENHOUSE AND INTERIORSCAPE PLANTS

For control of: Adelgids, Alder borer, Aphids, Asian longhorned beetle, Black vine weevil larvae, Bronze birch borer, Emerald ash borer, Eucalyptus longhorned borer, Flatheaded borers (including Bronze birch and Alder), Japanese beetles, Lace bugs, Leaf beetles (including elm and viburnum leaf beetles), Leafhoppers (including glassy-winged sharpshooter), Leafminers, Mealybugs, Pine tip moth larvae, Psyllids, Royal palm bugs, Sawfly larvae*, Soft scales, White grub larvae, Whiteflies
For suppression of: Armored scales, Thrips

USE RATE	APPLICATION SITE
TREES	
<p>Diameter at Breast Height (DBH) is measured at 4.5 feet from the ground. Use the following rates as a function of tree diameter at breast height (DBH): Apply 1 packet (1.6 oz) per 12 - 48 inches of trunk diameter (DBH). You may only use the higher rate on trees >15 inches (DBH) to control: Asian longhorned beetle, Emerald ash borer, Eucalyptus longhorned borer, Bronze birch borer, and Alder borer. To calculate the higher rates, divide trunk diameter by 12 - 23 inches. Refer to example calculations below. RESTRICTION: DO NOT apply more than 5.375 packets (8.6 oz) (0.4 lb AI) per acre per year.</p>	<p>Soil Injection: GRID SYSTEM: Holes must be spaced on 2.5 foot centers, in a grid pattern, extending to the drip line of the tree. CIRCLE SYSTEM: Apply in holes evenly spaced in circles, (use more than one circle dependent upon the size of the tree) beneath the drip line of the tree extending in from that line. BASAL SYSTEM: Space injection holes evenly around the base of the tree trunk no more than 6 to 12 inches out from the base. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per tree. NEW YORK SPECIFIC RESTRICTION: No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York. Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone. For Control of Specified Borers: Application to trees already heavily infested may not prevent the eventual loss of the trees due to existing pest damage and tree stress.</p>
<p>EXAMPLE CALCULATIONS: Example 1 (to calculate the standard rate): If you have three trees having DBH of 8, 10 & 16 inches, the total cumulative inches of trunk diameter is 34 inches (8 + 10 + 16 = 34) 34/48 = 0.708 x 1.6 fl. oz (1 packet) = 1.13 oz -OR- 34/24 = 1.417 x 1.6 fl. oz. (1 packet) = 2.27 oz If you have a single tree with a DBH of 12 inches, the lower rate range will be: 12/48 = 0.25 x 1.6 oz (1 packet) = 0.4 oz -OR- 12/24 = 0.5 x 1.6 oz (1 packet) = 0.8 oz Example 2 (To calculate the higher rate): If you have three trees having DBH of 15, 20 & 25 inches, the total cumulative inches of trunk diameter is 60 inches (15 + 20 + 25 = 60) 60/23 = 2.6 x 1.6 fl. oz (1 packet) = 4.17 oz -OR- 60/12 = 5 x 1.6 fl. oz. (1 packet) = 8 oz If you have a single tree with a DBH of 30 inches, the higher rate range will be: 30/23 = 1.3 x 1.6 oz (1 packet) = 2.08 oz -OR- 30/12 = 2.5 x 1.6 oz (1 packet) = 4.0 oz</p>	

SHRUBS

1 packet (1.6 oz) per 24 - 48 feet cumulative shrub height	Soil Injection: Apply to individual plants using dosage indicated. Mix required dosage in sufficient water to inject an equal amount of solution in each hole. Maintain a low pressure and use sufficient solution for distribution of the liquid into the treatment zone. Keep the treated area moist for 7 to 10 days. DO NOT use less than 4 holes per shrub. NEW YORK SPECIFIC RESTRICTION: No Soil Injection Applications Allowed in Nassau or Suffolk Counties of New York. Soil Drench: Uniformly apply the dosage in no less than 10 gallons of water per 1,000 square feet as a drench around the base of the tree, directed to the root zone. Remove plastic or any other barrier that will stop solution from reaching the root zone.
--	--

FLOWERS & GROUNDCOVERS

1 packet (1.6 oz) / 8,250 - 11,000 ft ² or 4 - 5.375 packets (6.4 - 8.5 oz) / Acre	Apply as a broadcast treatment and incorporate into the soil before planting or apply after plants are established. Irrigate immediately following application to established plants.
---	---

Remarks

*Sawfly larvae feed on mature foliage beginning in early spring. Make treatments in the fall before sawfly emergence in spring to allow adequate time for imidacloprid translocation into mature foliage.

Diameter at Breast Height (D.B.H.) is measured at 4.5 feet from the ground.

Restrictions

DO NOT apply more than 5.375 packets (8.6 oz) (0.4 lbs AI) per acre per year.

DO NOT harvest or consume fruits or nuts from trees that have been treated within 1 year of application.

FIELD AND FOREST NURSERIES

Pests	Rate	
For control of: White grub larvae ¹ (such as Japanese beetle, Masked chafers, European chafer, Oriental beetle, Asiatic garden beetle)	1 packet (1.6 oz) per 2500 ft of row	1 packet (1.6 oz) per 7000 ft ²

Application methods

Apply May through July. Time the treatment so that rainfall or irrigation occurs within 24 hours following the application.

Apply as a uniform band on either side of the row using a band width six (6) inches wider than the actual root ball diameter to be dug. DO NOT allow bands in adjacent rows to overlap.

Remarks

Mowing of the vegetation in the area to be treated to a height of 3 inches or less prior to application will improve the consistency of control.

Restrictions

DO NOT use less than 2 gallons of spray volume per 1,000 ft²

DO NOT apply more than 5.375 packets (8.6 oz) (0.4 lbs AI) per acre per year.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area. Exposure to moisture or excessive handling of water-soluble packets may cause breakage.

Pesticide Disposal: Pesticide spray mixture or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling: Nonrefillable container. DO NOT reuse or refill this container. Outer packaging for this product is secondary packaging to contain Water soluble packaging. Thoroughly rinse any soluble powder residue from container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC AT 1-800-424-9300

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of PrimeraTurf Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold PrimeraTurf Inc. and Seller harmless for any claims relating to such factors.

PrimeraTurf Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or PrimeraTurf Inc., and Buyer and User assume the risk of any such use. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW PrimeraTurf Inc. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

To the extent consistent with applicable law, PrimeraTurf Inc. or Seller shall not be held liable for any incidental, consequential or special damages resulting from the use or handling of this product. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF PrimeraTurf Inc. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF PrimeraTurf Inc. OR SELLER, THE REPLACEMENT OF THE PRODUCT.

PrimeraTurf Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of PrimeraTurf Inc.

PrimeraTurf and PrimeraOne are a trademarks of
PrimeraTurf, Inc.

Actara, Centric, and Platinum are registered
trademarks of the Syngenta Group Company.
Assail and Intruder are registered trademarks of
Nippon Soda Company, LTD.

Admire, Calypso, Leverage, and Trimax are
registered trademarks of Bayer CropScience.

Manufactured for:

PrimeraTurf Inc.

3677 Guildhall Trail

Marietta, GA 30066

**NET CONTENTS:
1.6 Ounces**