

BROADLEAF WEED KILLER with TRIMEC® HERBICIDE



CONTROLS: Dandelion, Chickweed, Knotweed, Plantain, Henbit, Spurge and many other broadleaf weeds, as listed on the back panel.

A	CT	IVI	E IN	GREDI	ENTS:
_					

2,4-dichlorophenoxyacetic acid	1.37%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid	0.31%
Dicamba: 3,6-dichloro-o-anisic acid	0.13%
OTHER INGREDIENTS:	98.19%
TOTAL	100.00%

THIS PRODUCT CONTAINS:

27.4 lb. 2,4-dichlorophenoxyacetic acid equivalent per ton.

6.2 lb. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.

2.6 lb. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods

EPA REG. NO. 2217-640-961

(L) (D) EPA Est. 961-PA-01; 961-IL-01 Superscript used is first letter of lot number.

TRIMEC[®] is a registered trademark of PBI/Gordon Corporation.

CAUTION

See back panel for complete Precautionary Statements and Directions for Use, including First Aid and Storage and Disposal.



Distributed by: Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042 1-800-233-0628 • (717) 273-1685 Manufactured in U.S.A.

NET WT. 20 LBS. (9.07 KG) Covers up to 10,000 sq. ft.



24-35799

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Causes moderate eye irritation. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Wear long-sleeved shirt and long pants, socks, shoes, and gloves. Remove and wash contaminated clothing before reuse.

	FIRST AID
If on skin or on clothing:	Take off contaminated clothing.
	Rinse skin immediately with plenty of water for 15 - 20 minutes.
	Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.
	Call a poison control center or doctor for further treatment advice.
If in eyes:	 Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.
	 Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
	 Call a poison control center or doctor for treatment advice.
If swallowed:	Call a poison control center or doctor immediately for treatment advice.
	Have person sip a glass of water if able to swallow.
	Do not induce vomiting unless told to do so by a poison control center or doctor.
	Do not give anything by mouth to an unconscious person.
Have the product conta	iner or label with you when calling a poison control center or doctor or going for treatment.
For additional informati	on in case of emergency, call toll free 1-877-800-5556.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS

- For residential use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.
- Apply when the air is calm. Do not apply to newly seeded lawns. Do not exceed specified dosages for any area. During application, close spreader openings when turning.
- Do not use this product on or near desirable plants, including within the drip line of the roots of desirable trees and shrubs, since injury may result.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

INSTRUCTIONS:

Lebanon Broadleaf Weed Killer with Trimec® Herbicide controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others as listed which are commonly found in home lawns. Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in 30 days.

For use on established turf of warm season grasses including bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses including Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum broadcast application rate is 2.0 pounds of product per 1,000 sq. ft. per application. The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications. The maximum seasonal rate is 4.0 pounds of product per 1,000 sq. ft.

- 1. Mow lawn to normal height 1 to 2 days before application.
- 2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
- 3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to allow the granules to adhere and to remain on the leaf surface of the weeds.
- 4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

SUGGESTED SPREADER SETTINGS								
Spreader	2#/1,000 sq. ft.	Spreader	2#/1,000 sq. ft.					
LebanonTurf	4	Lesco (letter/Numeric)	G/15					
Andersons AccuPro	Н	PennMulch HVO	J					
Earthway Rotary	16	ProScape SS	J					
Gandy	24	Spyker	4					
Lely	411	Vicon (all models)	22					
The above spreader settings are appro-	ximate. Calibrate your spreader on 10	00 square feet and adjust to apply at t	he rate of 1 pound of					
product per 500 square feet.	•	. , , , , , , , , , , , , , , , , , , ,	,					

Weeds

Wild four-o'clock Annual yellow Buckhorn Clover, white Hawkweed Mustard Prostrate vervain Spotted catsear Wild garlic Spotted spurge Buckhorn plantain sweetclover Cockle Healall Narrowleaf plantain Puncturevine Wild geranium Wild lettuce Heartleaf drymary Narrowleaf vetch Spurweed Bulbous buttercup Cocklebur Aster Purslane, common Stinging nettle Austrian fieldcress Bull thistle Common mullein Heath aster Nettle Ragweed Wild marigold Strawberry, India mock Bullnettle Hedge bindweed Orange hawkweed Red sorrel Bedstraw Creeping jenny Oriental cocklebur Tall nettle Wild mustard **Beggarticks** Burclover Cudweed Hedge mustard Redroot pigweed Wild onion Tall vervain Betony, Florida Burdock Curly dock Hemp Oxalis Redstem filaree Tansy ragwort Wild parsnip Bindweed, field Burning nettle Daisy, English Parsley-piert Henbit Rough cinquefoil Daisy fleabane Tansy mustard Wild radish Burweed Hoary cress Parsnip Bird vetch Rough fleabane Wild rape Tanweed Bitter wintercress Buttercup Hoary plantain Daisy, oxeye Pearlwort Russian pigweed Wild strawberry Thistle Buttonweed Dandelion Bittercress, hairy Hoary vervain Pennycress Russian thistle Bitterweed Canada thistle Dichondra Pepperweed Trailing crownvetch Wild sweet potato Horsenettle Scarlet pimpernel Wild vetch Tumble mustard Black-eyed Susan Carolina geranium Dogbane Pigweed Scotch thistle Jimsonweed Woodsorrel Tumble pigweed Black medic Carpetweed Pineywoods bedstraw Dogfennel Knawel Sheep sorrel Dollarweed Velvetleaf Woolly croton Black mustard Catchweed bedstraw Knotweed Plains coreopsis Shepherdspurse Woolly morningglory Venice mallow Blackseed plantain Catnip False dandelion Kochia Plantain Slender plantain Woolly plantain Virginia buttonweed Blessed thistle Catsear Falseflax Lambsquarters Poison ivy Smallflower galinsoga Virginia creeper Wormseed Bloodflower milkweed Chickweed, common False sunflower Poison oák Smooth dock Lespedeza Virginia pepperweed Yarrow Chickweed, mouseear Mallow Pokeweed Smooth pigweed Blue lettuce Fiddleneck Wavyleaf bullthistle Yellow rocket Chicory Florida pusley Blue vervain Matchweed Poorjoe Sorrel Western clematis Yellowflower Bracted plantain Cinquefoil Frenchweed Mexicanweed Prairie sunflower Sowthistle Western salsify pepperweed Clover, crimson Prickly lettuce Spanishneedles Brassbuttons Galinsoga Milk vetch White mustard Clover, hop Prickly sida Bristly oxtongue Goldenrod Morningglory Speedwell Mouseear hawkweed Wild aster Broadleaf dock Clover, red Prostrate knotweed Spiny amaranth Ground ivy Wild buckwheat Broadleaf plantain Clover, strawberry Gumweed Prostrate pigweed Spiny cocklebur Mugwort Wild carrot Clover, sweet Hairy fleabane Musk thistle Prostrate spurge Broomweed Spiny sowthistle

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING:

Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER.

FOR USE ONLY AS DIRECTED.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.