

Net Wt. 50 Lbs. (22.68 kg)

Turf Products Fertilizer with 0.86% ProPendi™ Herbicide 20-0-4

Tall fescue

 Contains Fortify® -N KEEP OUT OF REACH OF CHILDREN PRECAUTIONARY STATEMENTS

.375 Seal

Hazards to Human and Domestic Animals: CAUTION. May be harmful if swallowed. Harmful if absorbed through the skin. Causes moderate eye irritation. Avoid contact with eyes, skin, or clothing. If swallowed: Call a poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce DIRECTIONS FOR USE

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains

center or doctor, or going for treatment. You may also contact 1-800-757-8951 for emergency medical treatment information.

Personal Protective Equipment (PPE)

Follow manufacturer's instructions for cleaning/maintaining personal detergent and hot water. Keep and wash personal protective equipment greenhouses. separately from other laundry.

 Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. thoroughly and put on clean doming.

Remove PPE immediately after handling this product. Wash the outside

Liability" statements.

Tall Fescue purslane

(WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS. either collect for use or dispose of properly. ACTIVE INGREDIENT:

Underlined letter used is first letter of run code on end of bag. Guaranteed Analysis 20.00% Urea Nitrogen* Derived from urea and potassium chloride *6.00% nitrogen stabilized with N-(n-butyl) thiophosphoric triamide

Information regarding the contents and levels of metals in this product is available on the internet at http://www.regulatory-info-ap.com We recommend that you follow the Florida Green Industries BMP's at: http://fyn.ifas.ufl.edu/pdf/GIBMP_Manual_Web_English_2015.pdf

controlled-release nitrogen feeding: Grassy Weeds Dicot Weeds crabgrass chickweed goosegrass knotweed Poa annua prostrate spurge barnyardgrass purslane

vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin

With its labeling.

Do not apply this product in a way that will contact workers or other

Comments immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

It is proved that will contact with earlier and with plenty of water for 15-20 minutes. Call a poison control the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth,

an ambulance of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and if possible. Call a poison control center or doctor for further treatment advice.

Contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry-interval. The requirements in this box only apply to uses of the product that are covered by the Worker Protection

Have the product container or label with you when calling poison control only apply to uses of the product that are covered by the Worker Protection

Standard.

Totalination, notification and exceptions and exceptions and exceptions and exceptions to weed germination to weed germination only apply to uses of the product that are covered by the Worker Protection

Not for use on greens and barryardgrass

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Some of the materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart. Applicators and other handlers must wear: Long-sleeved shirt and long pants, chemical resistant resistant gloves made of any waterproof material, shoes plus socks.

is used to produce agricultural plants on farms, forests, nurseries, or

 Wash hands before eating, drinking, chewing gum, using tobacco or
 For use on established turfgrasses including bentgrass, bermudagrass, and
 St. Augustinegrass
 Poa annua Poa annua (except putting greens and tees), Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass, and blends of the grasses. See "Bermudagrass, Bentgrass and Poa Annua Precautions" and "Limitation of Liability" statements.

Bahiagrass barnyardgrass Zoysiagrass fall panicum oxalis centipedegrass, and blends of the grasses. See "Bermudagrass, Bentgrass and Poa Annua Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass, Poa annua and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and Poa annua are the desired species, a light-rate application only is recommended.

Not for use on bentgrass, Poa annua and bermudagrass putting greens and tees)

(Not for use on greens and tees)

Comments How/When to Use
For control of annual grassy and broadleaf weeds such as crabgrass,

— control or where heavy weed infestations are expected.

goosegrass, foxtail, barnyardgrass, fall panicum, Poa annua, oxalis. knotweed, purslane, and prostrate spurge, apply in the spring before they Environmental Hazards: This pesticide is toxic to fish. Do not apply sprout.

Comments areas below the mean high water mark. Runoff from treated areas may be

To prevent Poa annua and fall germinating chickweed, apply product to dry

Apply a repeat application of 175-233 lbs./acre to extend control of gooseg
burned, stay out of smoke. hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, aither collect to use of proposity.

IMPORTANT: READ BEFORE USE: Read the entire Directions for Use and the activity of the lower rate was used initially.

IMPORTANT: READ BEFORE USE: Read the entire Directions for Use and the entire D

Where lower infestations are expected, the lighter rates may be used. 9198-AL-001A when repeat applications are necessary after 6-8 weeks.

To provide proper distribution calibrate spreader before application. Do not adjust rates in order to meet fertilizer requirements of your soil. 50 lbs. treats 15,000 sq. ft. (0.34 acre) at the 145 lbs. product/acre - LIGHT RATE (3.32 lbs./1,000 sq. ft.) 50 lbs. treats 12,400 sq. ft. (0.28 acre) at the 175 lbs. product/acre - LOW RATE (4.01 lbs./1,000 sq. ft.) 50 lbs. treats 10,700 sq. ft. (0.25 acre) at the 203 lbs. product/acre - MEDIUM RATE (4.66 lbs./1,000 sq. ft.) 50 lbs. treats 9,300 sq. ft. (0.21 acre) at the 233 lbs. product/acre - HIGH RATE (5.35 lbs./1,000 sq. ft.)

Prevents sprouting of the following grassy and dicot weeds and provides a Do not mow treated area until product has been watered in and turf foliage Turfgrass Species Weeds Rates Cool Season Grasses Crabgrass 175-233 lbs./acre as the initial

Perennial ryegrass Poa annua germination Fine fescue barnyardgrass fall panicum prostrate spurge

Kentucky bluegrass foxtail application prior to weed

prior to weed germination. Apply a repeat application after 6-8 weeks at 175-203

prostrate spurge knotweed

handlers must wear: Long-sleeved shirt and long pants, chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to use of this product that are not within the scope of the Worker Protective Standard for agricultural pesticides (40)

Goosegrass 175 lbs./acre in the spring prior to weed germination Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dusts

Warm Season Grasses crabgrass 175-233 lbs./acre prior to have settled.

| Have settled | Have

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard

That meets the requirements listed in the Worker Protection Standard

The worke

weed germination cudweed 175-233 lbs./acre prior to Read the entire Directions for Use and the Warranty Disclaimer and hop clover weed germination

eveningprimrose

Pendimethalin † 0.86% Apply product uniformly to dry foliage at the suggested rate with a properly INERT INGREDIENTS: 99.14% calibrated spreader. For best results, water after application (1/4-1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier of the surface for an effective chemical barrier. Do not destroy the chemical barrier of the surface for an effective chemical barrier. Do not destroy the chemical barrier of the surface for an effective chemical barrier. Do not destroy the chemical barrier of the surface for an effective chemical barrier. Do not destroy the chemical barrier of the surface for an effective chemical barrier. †CAS # 40487-42-1 created at the soil surface by raking or aerifying as a segment to rate was used initially.

EPA Reg. No. 9198-173 EPA Est. No. 9198-OH-1M, 9198-OH-2B, application. Allow at least a 4 month space between the applications except GROUND WIDTH OF LIGHT RATE LOW RATE MEDIUM RATE HIGH RATE
 SPEED
 COVERAGE
 1.24 lbs.ai/acres
 1.50 lbs. ai/acres
 1.75 lbs. ai/acres
 2.00 lbs. ai/acres

Do not use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa*

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's. This product may cause discoloration and thinning where over applications have been made if the turfgrass is under serve stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil

Do not apply to dormant bent-grass and *Poa annua* (turf that is not actively

may be extensive in low-cut bermudagrass and on sandy soils. Goosegrass 175-233 lbs./acre in the spring As with most other preemergence herbicides, application of this product to bermudagrass overseeded with cool season turfgrass species may injure Do not apply to bentgrass or *Poa annua* after late summer.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse of with water. at the application site, do not use this product under windy conditions (5-10 mph) due to its staining potential. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not

feed clippings to livestock. Properties of All Dinitroanaline (DNA) Products May cause temporary slowing of turf fill-in and stolon-rhizome rooting

Requires a waiting period before seeding or sprigging. Seeding or Sprigging Intervals after Treatment surface should be disturbed or worked to dilute any remaining chemical

Treatment Interval after Seeding or Sprigging
On newly planted areas, wait until the grass has been mowed at least 4

Do not apply more than a cumulative total of 349 lbs. of product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 698 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year on cool season year on warm season turfgrasses. STORAGE AND DISPOSAL Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a clean, dry place. Reseal opened bag by

Goosegrass 175-233 lbs./acre prior to

disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this by incineration, or, if allowed by State and local authorities, by burning. If

WARRANTY DISCLAIMER and LIMITATION of LIABILITY: ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS; YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE/JURISDICTION P.O. Box 119, Maumee, OH 43537



Andersons®

Visit our website at www.AndersonsPlantNutrient.com

©2019 The Andersons, Inc. The Andersons logo is a registered trademarks of The Andersons, Inc.; ProPendi is a trademark of The Andersons, Inc. Fortify-N is a registered trademark of The Andersons, Inc.

> **APT20UPD5.40** APT20UPD5.40B

Turf Products Fertilizer with

Andersons 0.86% ProPendi™ Herbicide 20-0-4

SEAL AREA NO COPY

SEAL AREA NO COPY

SEAL AREA NO COPY

SEAL AREA NO COPY